

A Fluid Generalization of Membranes.

Mark D. Roberts, University of Surrey, GU2 7XH,
max1mr@eim.surrey.ac.uk, <http://cosmology.mth.uct.ac.za/~roberts>

February 1, 2008

Abstract

In a certain sense a perfect fluid is a generalization of a point particle. This leads to the question as to what is the corresponding generalization for extended objects. The lagrangian formulation of a perfect fluid is much generalized and this has as a particular example a fluid which is a classical generalization of a membrane, however there is as yet no indication of any relationship between their quantum theories.

Contents

1	Introduction.	1
2	The Perfect Fluid.	3
3	The Dirac Membrane.	5
4	The F-fluid.	6
5	The F-brane.	7
6	Conclusion.	8

1 Introduction.

There are three main types of dynamical system which have a lagrangian formulation: fields theories, extended objects, and fluids. Some other systems which have lagrangian formulation include relative motion systems [16]. Various relationships have been found between string theories and field theories [6]p.26-27, also various string theories can be written as field theories [17]. Some relationships are known between extended objects and fluids [10][9][5], these seem to be of the form of extended objects producing fluids. In the present work it is fluids which are the more general object and the fluids reduce to extended objects. A generalization of string theory called M-theory is currently being sought, whether there is any relationship between

this hypothetical generalization and fluids remains to be seen. The prospect of generalizing the lagrangian theory of fluids rests on the observations that a perfect fluid spacetime stress can be reduced to that of a congruence of point particles. It seems that imperfect fluids do not often have a lagrangian description, in particular heat conduction and anisotropic stress do not seem to be derivable from a lagrangian. The lagrangian theory of fluids involves all three of: relativity, quantum theory and thermodynamics, although there has not yet been any connection established with quantum field theory on curved spacetime [1] where this is also the case.

There are at least **four** other reasons for studying lagrangian fluid theory.

- symmetry breaking, see [13],
- low temperature physics, so far there has been no contact at all of Lagrangian based theory with existing approaches, such as [3],
- quantum cosmology, with the geometry coupled to a fluid, see for example [11],
- the study of irreversible process, see for example [8], a lagrangian form might give insight into the processes involved.

A perfect fluid has a gauge description [14]; this allows it to be canonically quantized producing a quantum fluid theory which has novel quantum algebra, see [14] and §5.

In §2 the properties of a perfect fluid are described. In §3 some of the properties of Dirac membranes are described, the treatment of the constraints here is a different from usual. In §4 the lagrangian description of a perfect fluid is much generalized, this involves putting internal indices on fluid objects, the indices label the object and do not usually involve differentiation. Up to adding similarly structured terms, it is hoped that the resulting f -fluid is the most general that can be derived by a spin-free lagrange method; however it is not general enough to incorporate heat conduction or anisotropic stresses. In §5 it is shown how to reduce the number of indices by using an internal metric, this produces the f -brane for which both lagrangian and metric stress have both the perfect fluid and membranes as examples. Reduction is taken to have happened when both the lagrangian and metric stress coincide. The notation used is: signature $(-+++)$, greek indices $\alpha, \beta, \dots, \mu, \dots = 0, 1, 2, 3$ are spacetime indices, early latin indices $a, b, c, \dots$ are internal or fluid indices, middle greek indices $\iota \dots$ index constraints if they are not otherwise indexed, middle latin indices $i, j, k, \dots$ are velocity potential indices, all indices are left out when it is hoped that the ellipsis is clear, γ is the auxiliary metric which is usually that of the internal space of a membrane, $\wp$ is the pressure but $p + 1$ the dimensions of γ , μ is the density, n is the particle number, ξ is the enthalpy but h is the trace of the projection tensor and $\hbar$ is Planck's constant divided by 2π , W is a vector field which has been decomposed into clebsch velocity potentials, V is a timelike vector field constructed from W and the enthalpy ξ . All other conventions are those of Hawking and Ellis [7].

2 The Perfect Fluid.

The lagrangian of a perfect fluid is taken to be the pressure

$$\mathcal{L} = \wp, \quad (1)$$

and the hamiltonian is the density μ . The lagrangian 1 is varied via the first law of thermodynamics

$$d\wp = nd\xi - nTds, \quad (2)$$

where n is the particle number, ξ is the enthalpy, T is the temperature, and s is the entropy. The fluid velocity vector V is a unit timelike vector field which has a clebsch decomposition

$$\xi V_\mu = W_\mu = \sigma_\mu + \theta s_\mu + \dots, \quad V^\mu V_\mu = -1, \quad h_{\mu\nu} = g_{\mu\nu} + V_\mu V_\nu, \quad (3)$$

where σ, θ, s are the clebsch potentials, σ is the higgs [13], θ is the thermasy, and s the entropy. The number of extra terms depends on the dimension of the spacetime involved. Clebsch's theorem is only local, so that when there are obstructions to the velocity V or unusual global properties it no longer holds; the usual way around this is to add yet more terms to the decomposition. For our purposes when the first three terms are known it is straightforward to add the extra terms, so that we use just three terms regardless of dimension and so on. Using the unit normalization of V , $V^2 = -1$ and $\xi V = W$ the first law 2 can be written

$$d\wp = -nV_\mu dW^\mu - nTds, \quad (4)$$

using

$$\wp + \mu = n\xi, \quad (5)$$

and varying with respect to the metric gives the metric stress

$$T_{\mu\nu} = (\wp + \mu)V_\mu V_\nu + \wp g_{\mu\nu}. \quad (6)$$

The reduction to a congruence of point particles is achieved using

$$\mathcal{L} = \wp = -m\ell = -m\sqrt{-\dot{x}^2}, \quad \xi = \pm\ell, n = \pm m, \quad \mathcal{H} = \mu = 0, \quad V^\mu = \frac{\dot{x}^\mu}{\ell}, \quad T_{\mu\nu} = m\ell h_{\mu\nu}. \quad (7)$$

The point particles equation of motion is $\dot{P}/m = \dot{V} = 0$ is not explicitly recovered, $\dot{V}^a = V_b V_b^a$ is just the acceleration of the fluid, so that in addition to 7 the reduction requires that the fluid is acceleration free. The reduction to a single point particle is achieved by either using delta functions or by varying the lagrangian with respect to the single internal index, see [15]. Variation of 1 with respect to the clebsch potentials give their equations of motion

$$(nV^\mu)_{;\mu} = \dot{n} + n\Theta = 0, \quad \dot{s} = 0, \quad \dot{\theta} = T. \quad (8)$$

The momenta are

$$\Pi^\sigma = -n, \quad \Pi^\theta = 0, \quad \Pi^s = -n\theta, \quad (9)$$

and the constraints between the momenta are

$$\phi_1 \equiv \Pi^s - \theta \Pi^\sigma, \quad \phi_2 \equiv \Pi^\theta. \quad (10)$$

The Poisson bracket is defined by

$$\{A, B\} \equiv \frac{\delta A}{\delta q_i} \frac{\delta B}{\delta \Pi^i} - \frac{\delta B}{\delta q_i} \frac{\delta A}{\delta \Pi^i}. \quad (11)$$

The Dirac matrix is defined by

$$C_{\iota\kappa} \equiv \{\phi_\iota, \phi_\kappa\}, \quad (12)$$

and the Dirac bracket is defined by

$$\{A, B\}^* = \{A, B\} - \{A, \phi_\iota\} C_{\iota\kappa}^{-1} \{\phi_\kappa, B\}. \quad (13)$$

Quantization is done using the substitution of the Dirac bracket by commutators

$$\{A, B\}^* \rightarrow \frac{1}{i\hbar} [\hat{A}\hat{B} - \hat{B}\hat{A}]. \quad (14)$$

For the perfect fluid

$$C_{12} = \{\phi_1, \phi_2\} = \Pi^\sigma, \quad C_{\iota\kappa} = -i\sigma^2 \Pi^\sigma, \quad C_{\iota\kappa}^{-1} = \frac{i\sigma^2}{\Pi^\sigma}, \quad (15)$$

where σ^2 is the Pauli matrix

$$\sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}. \quad (16)$$

Forming the Dirac brackets between the fields and momenta it turns out that it is possible to multiply by Π^σ throughout. After applying the quantization 14 and dropping the hats the equations between the fields and momenta are

$$\begin{aligned} [\sigma\theta - \theta\sigma]\Pi^\sigma &= \Pi^\sigma[\sigma\theta - \theta\sigma] = -i\hbar\theta\delta^4(x-y), [\theta s - s\theta]\Pi^\sigma = \Pi^\sigma[\theta s - s\theta] = -i\hbar\delta^4(x-y), \\ [\Pi^\sigma\sigma - \sigma\Pi^\sigma] &= -i\hbar\delta^4(x-y), [\Pi^s - s\Pi^s] = -i\hbar\delta^4(x-y). \end{aligned} \quad (17)$$

Using the substitutions

$$v_1 = \sigma, \quad v_2 = s, \quad v_3 = \theta, \quad v_4 = \Pi^\sigma, \quad v_5 = \Pi^s, \quad v_6 = \Pi^\theta, \quad (18)$$

setting $\hbar = 1$, and then suppressing the delta function $\delta^4(x-y)$ gives

$$v_4(v_3v_2 - v_2v_3) = -i, \quad v_4(v_1v_3 - v_3v_1) = -iv_3, \quad v_4v_1 - v_1v_4 = -i, \quad v_5v_2 - v_2v_5 = -i. \quad (19)$$

v_6 does not occur. In the case when V is a gradient vector $V^\mu = \sigma^\mu$, only the second to last of these commutators remains, so that the algebra is the same as that of the point particle.

3 The Dirac Membrane.

The lagrangian is [4]

$$\mathcal{L} = k\sqrt{-\gamma}, \quad \gamma_{ab} = x_{\mu a}x_b^\mu, \quad \sqrt{-\gamma} = (-\det \gamma_{ab})^{\frac{1}{2}}. \quad (20)$$

The first fundamental form is

$$\aleph^{\mu\nu} = \gamma^{ab}x_a^\mu x_b^\nu, \quad \aleph^{\mu\rho}\aleph_\rho^\nu = \aleph^{\mu\nu}, \quad \aleph_\rho^\rho = \gamma_c^c = x^{\rho c}x_{\rho c} = p + 1. \quad (21)$$

The projection tensor is

$$h^{\mu\nu} \equiv g^{\mu\nu} - \aleph^{\mu\nu}, \quad h = d - 1 - p. \quad (22)$$

The metric stress is

$$T^{\mu\nu} = k\sqrt{-\gamma}(-(p+1)\aleph^{\mu\nu} + g^{\mu\nu}). \quad (23)$$

The momenta are

$$P_\mu^a = -k\sqrt{-\gamma}x_\mu^a. \quad (24)$$

The constraints between the momenta are

$$\phi_{\mu\nu}^{ab} \equiv P_\mu^a P_\nu^b + k^2 \gamma x_\mu^a x_\nu^b, \quad (25)$$

compare [12] and [2]. 25 has the contractions

$$\begin{aligned} \phi_{\mu\nu} = \phi_{a\mu\nu}^a &= P_\mu^a P_{\nu a} + k^2 \gamma \aleph_{\mu\nu}, \quad \phi^{ab} = \phi^{ab\mu}_\mu = P^{a\mu} P_\mu^b + k^2 \gamma \gamma^{ab}, \\ \phi &= \phi_\mu^\mu = \phi_a^a = P^{a\mu} P_{a\mu} + k^2 (p+1) \gamma. \end{aligned} \quad (26)$$

Using just the last of these it is possible to derive a Klein-Gordon equation with $k^2 = m^2$, see [16]; however in general one needs the other constraints. Define

$$X_c^{ab\mu} \equiv \frac{\delta \phi^{ab}}{\delta P^{c\mu}} = 2\delta_c^{(a} P^{b)\mu} \quad (27)$$

$$Y_\mu^{ab} \equiv \frac{\delta \phi^{ab}}{\delta x^\mu} = -2k^2(\gamma x_\mu^{(a})^b - 2k^2(\gamma \gamma^{ab} x_\mu^c)_c = -2k \left(\sqrt{-\gamma}(P_\mu^c \gamma^{ab} + P_\mu^{(a} \gamma^{b)c}) \right)_c.$$

X has one more index than Y , so that it is not possible to form a Dirac matrix C for them as they are. Choosing $c = \tau$ in X one can form $C_{abcd} = \{\phi_{ab}, \phi_{cd}\}$, but it is not clear what the inverse of this should be. Choosing $b = c = \tau$ in X and $b = \tau$ in Y one sees that the constraints are limited to the $(p+1)$ first class constraints $\phi^a = \phi^{a\tau}$ and one can form Dirac matrix $C_{ab} = \{\phi_a, \phi_b\}$. Restricting to the case of the string $p = 1$ and

$$C_{ab} = i\sigma^2 C_{\tau\sigma} = i\sigma^2 Z^{\tau\sigma\mu}_\mu, \quad C_{ab}^{-1} = \frac{-i\sigma^2}{Z^{\tau\sigma\mu}_\mu}, \quad Z^{\tau\sigma}_{\mu\nu} \equiv Y_\nu^{\tau\tau} X_{\tau\mu}^{\sigma\tau} - Y_\nu^{\sigma\tau} X_{\tau\mu}^{\tau\tau}. \quad (28)$$

The Dirac brackets between the coordinates and momenta are

$$\{x^\mu, x^\nu\}^* = \{P^\mu, P^\nu\}^* = 0, \quad \{x^\mu, P^\nu\}^* = g^{\mu\nu} - i\sigma^2 \frac{Z^{\tau\sigma\mu\nu}}{Z^{\tau\sigma\rho}_\rho}. \quad (29)$$

One can get an explicit form of these Dirac brackets by using a specific form of the internal metric γ , say the Nambu-Goto choice

$$\gamma_{ab} = \begin{pmatrix} \dot{x}^2 & \dot{x} \cdot x' \\ x' \cdot \dot{x} & x'^2 \end{pmatrix}, \quad \gamma\gamma^{ab} = \begin{pmatrix} x'^2 & -x' \cdot \dot{x} \\ -\dot{x} \cdot x' & \dot{x}^2 \end{pmatrix}, \quad \gamma = -\mathcal{A}^2 = -\dot{x}^2 x'^2 + (\dot{x} \cdot x')^2, \\ k = \frac{1}{2\pi\alpha'}, \quad 2\pi\alpha' \mathcal{A}P^{\tau\mu} = x'^2 \dot{x}^\mu - \dot{x} \cdot x' x'^\mu, \quad 2\pi\alpha' \mathcal{A}P^{\sigma\mu} = \dot{x}^2 x'^\mu - \dot{x} \cdot x' \dot{x}^\mu, \quad (30)$$

however from the present perspective the important point is that now the $\{x, P\}^*$ Dirac bracket is different from the Poisson bracket, whereas for the perfect fluid the $\{q, q\}^*$ bracket also differs.

4 The F-fluid.

The lagrangian is taken to be

$$\mathcal{L} = f(\wp_1, \wp_2, \dots, \wp_F), \quad (31)$$

the pressures $\wp$ and densities μ are equated to the enthalpies ξ and particle numbers n by

$$\wp_a + \mu_a = n_a^{bc} \xi_{bc}, \quad (32)$$

the internal indices $a, b, \dots$ label distinct objects, for example $\wp_1, \wp_2, \dots$ and only indicate differentiation with respect to the index when it is on x or is outside a bracket $()_a$. The unit timelike condition on V generalizes to

$$h_b^c V_\mu^b = W_\mu^c, \quad V_d^\mu V_\mu^e = -\delta_d^e, \quad (33)$$

so that

$$-\xi_{ab} = -\delta_b^c \xi_{ac} = \xi_{ac} V_b^\mu V_\mu^c = V_b^\mu W_{a\mu}. \quad (34)$$

This allows the second equality in the first law

$$dp_a = n_a^{bc} d\xi_{bc} - n_a^{bc} T_b ds_c = -n_a^{bc} V_{\mu b} dW_\mu^c - n_a^{bc} T_b ds_c. \quad (35)$$

The metric stress is

$$T_{\mu\nu} = f_{,\wp_a} n_a^{bc} \xi_{ce} V_{\mu b} V_\nu^e + f g_{\mu\nu}. \quad (36)$$

The decomposition of W is

$$W_\mu^c = \sigma_\mu^c + \alpha_{ab}^c \theta^a s_\mu^b + \dots \quad (37)$$

In the present case this decomposition is an assumption, unlike for the perfect fluid where it is a consequence of Clebsch's theorem, α_{ab}^c is a new object. Varying with respect to the potential gives the equations of motion

$$(V_b^\mu n_a^{bc})_\mu = 0, \quad n_a^{bc} \alpha_{cde} V_b^\mu s_\mu^e = 0, \quad V_b^\mu n_a^{bc} (\alpha_{cde} \theta^d)_\mu - n_a^{be} T_b = 0, \quad (38)$$

for σ^a , θ^a , s^a respectively. The momenta are

$$\Pi_a^{\sigma bc} = -n_a^{bc}, \quad \Pi_a^\theta = 0, \quad \pi_a^{sbc} = -n_a^{bc} \alpha \cdot \theta, \quad (39)$$

where $\alpha_a = \alpha_{ac}^c$. The constraints are

$$\phi_a^{bc} \equiv \Pi_a^{sbc} - \alpha \cdot \theta \Pi_a^{\sigma bc}, \quad \phi_a^2 \equiv \Pi_a^\theta. \quad (40)$$

In this form it is not possible to get an inverse Dirac matrix.

5 The F-brane.

The f -brane is a particular case of the f -fluid. First reduce the number of indices by

$$n_a^{bc} = \gamma^{bc} n_a, \quad \xi_{bc} = \frac{1}{p+1} \xi \gamma_{bc}, \quad (41)$$

where γ is an internal metric, this gives

$$\wp_a + \mu_a = \xi n_a, \quad \xi V_\mu^c = W_\mu^c, \quad V_\mu^c V_e^\mu = -\delta_e^c. \quad (42)$$

The first law is

$$dp_a = n_a \gamma^{bc} d(\xi \gamma_{bc}) - \gamma^{nc} n_a T_b ds_c = -n_a V_\mu^c dW_c^\mu - n_a T^c ds_c, \quad (43)$$

where one can take γ through the d to get the second equality. The metric stress is

$$T_{\mu\nu} = f_{,\wp_a} n_a \xi V_\mu^c V_{c\nu} + f g_{\mu\nu}. \quad (44)$$

The best identification of V is not immediate. The problem is that for the point particle the explicit normalization condition is $V_{\mu\tau} V_\tau^\mu = -1$, with τ subscripted both times, for only one internal index this does not matter, but for more than one internal index one wants to preserve covariance, so that a different choice has to be made, choosing

$$V^{\mu c} = \beta x^{\mu c}, \quad (45)$$

the normalization condition 41 and the definition of γ 41 require

$$V_\mu^c V_e^\mu = \beta^2 x^{\mu c} x_{\mu e} = \beta^2 \gamma_e^c = -\delta_e^c \Rightarrow \beta = i. \quad (46)$$

With this choice it is possible to introduce the first fundamental form into the metric stress

$$T_{\mu\nu} = -f_{,\wp_a} n_a \xi \aleph_{\mu\nu} + f g_{\mu\nu}, \quad (47)$$

choosing the simplest lagrangian

$$\mathcal{L} = f = \Sigma \wp_a, \quad (48)$$

gives the metric stress of the f -brane

$$T_{\mu\nu} = -\Sigma(\wp_a + \mu_a) \aleph_{\mu\nu} + \Sigma \wp_a g_{\mu\nu}. \quad (49)$$

This is much as expected, the only other possibilities for its form would be terms summing the internal indices occurring across the pressure part and the x part. The f -brane reduces to the Dirac membrane when

$$\Sigma \wp_a = k \sqrt{-\gamma}, \quad \Sigma \mu_a = p k \sqrt{-\gamma}, \quad (50)$$

different choices can reduce to the stress of the conformal membrane [15]§4. Varying with respect to the potentials the equations of motion are

$$(V^{\mu c} n_a) = 0, \quad n_a \alpha_{de}^b V_b^\mu s_\mu^e = 0, \quad V^{\mu c} (\alpha_{cde} \theta^d)_\mu - T^e = 0. \quad (51)$$

The momenta are as for the f -fluid, however choosing $\Pi_a^\sigma \gamma^{bc} = \Pi_a^{\sigma bc}$ and similarly for Π^s gives

$$\Pi_a^\sigma = -n_a, \quad \Pi_a^\theta = 0, \quad \Pi_a^s = -\alpha \cdot \theta n_a, \quad (52)$$

which have constraints

$$\phi_a^1 \equiv \Pi_a^s - \alpha \cdot \theta \Pi_a^\sigma, \quad \phi_a^2 \equiv \Pi_a^\theta. \quad (53)$$

These constraints give Dirac bracket

$$C_{ab}^{12} = \{\phi_a^1, \phi_b^2\} = \alpha_b \Pi_a^\sigma. \quad (54)$$

To proceed it is necessary to find an inverse of this. The simplest way is to reduce it from a four indexed matrix to a two index matrix by tracing across the internal indices

$$C_a^{12} = \{\phi_a^1, \phi_a^2\} = \alpha \cdot \Pi^\sigma, \quad (55)$$

and then proceed as for the perfect fluid in §2. Defining v 's as in 18, except that this time they all are subscripted by an internal index, the following algebra is found

$$\begin{aligned} \alpha \cdot v^4(v_a^1 v_b^3 - v_b^3 v_a^1) &= -i\alpha \cdot v^3 \delta_{ab}, & \alpha \cdot v^4(v_a^3 v_b^2 - v_b^2 v_a^3) &= -i\delta_{ab}, \\ \alpha \cdot v^4(v_a^3 v_b^6 - v_b^6 v_a^3) &= -i(\alpha \cdot v^4 \delta_{ab} - \alpha_b v_a^4), & v_a^4 v_b^1 - v_b^1 v_a^4 &= -i\delta_{ab}, & v_a^5 v_b^2 - v_b^2 v_a^5 &= -i\delta_{ab}. \end{aligned} \quad (56)$$

which reduces to 19 when $a, b, \dots = 1$. Note that v^6 occurs in 57 but not in 19.

6 Conclusion.

A classical perfect fluid can be reduced to a congruence of point particles using 7. The perfect fluid has one propagating degree of freedom, the particle number, and two independent functions in the stress, the pressure and density; whereas the congruence of point particles has one propagating degree of freedom, and one independent function in the stress. The corresponding quantum theory reduces when the velocity vector W is a gradient vector. In the classical case a perfect fluid can be generalized to a f -brane 48,49 which reduces to the Dirac membrane. The f -brane has $p+1$ propagating degrees of freedom, the $p+1$ particle numbers, and $2p+2$ independent functions in the stress. There is no indication that this correspondence remains so quantum mechanically, this is because: fluid momenta lack a spacetime index, whereas membrane momenta have a spacetime index; for membranes it is the $\{\Pi, x\}$ Dirac bracket which differs from the Poisson bracket, whereas for fluids the $\{q, q\}$ brackets differs as well.

References

- [1] Birrel, N.D. & Davies, P.C.W.
Quantum Fields in Curved Space.
Cambridge University Press (1982).

- [2] Cartas-Fuentevilla,R.
Indentically closed two-form for covariant phase space quantization of the Dirac-Nambu-Goto p-brane in a curved spacetime.
`hep-th/0204133` *Phys.Lett.***B536**(2002)283-288.
- [3] Carter,B & Khalatnikov,I.M.
Momentum, vorticity, and helicity in covariant superfluid dynamics.
*Math.Rev.93m:83046 Ann.Physics***219**(1992)243-265.
- [4] Dirac,P.A.M.(1962)
An Extendible Model of the Electron.
*Proc.Roy.Soc.***A268**,57-67.
- [5] Gibbons,G, Hori, Kentaro, Yi, Piljin
String Fluid from Unstable D-branes.
*Nucl.Phys.***B596**(2001)136-150. `hep-th/0009061`
- [6] Green,M.B., Schwarz,J.H. & Witten,E.
Superstring Theory: Volume 1 Introduction.
Cambridge University Press (1987).
- [7] Hawking, S. W. and Ellis, G. F. R. (1973)
The Large Scale Structure of Space-time.
*Math.Rev.***54** #12154, Cambridge University Press.
- [8] Hiscock,W.A. and Lindblom,L.
*Phys.Rev.***D 31**(1985)725.
- [9] Hoppe, Jens
Surface Motions and Fluid Dynamics.
`hep-th/9405001` *Phys.Lett.***B335**(1994)41-44.
- [10] Jackiw,R.
A Particle Field Theorist's Lectures on Supersymmetric, Non-Abelian Fluid Mechanics and d-Branes.
`hep-th/0010042`
- [11] Lapchinskii,V.G. & Rubakov,V.A.
*Theor.Math.Phys.***33**(1977)1076.
- [12] Pavšič, Matej
Phase Space Action for Minimal Surfaces of any Dimension in Curved Space-time.
*Math.Rev.89b:83016 Phys.Lett.***B197**(1987)327-331.
- [13] Roberts,M.D.
Fluid Symmetry Breaking II: Velocity Potential Method.
*Hadronic J.***20**(1997)73-84. `hep-th/9904079`

- [14] Roberts,M.D.
The Quantum Commutator Algebra of a Perfect Fluid.
*Mathematical Physics, Analysis and Geometry***1**(1999)367-373. [gr-qc/9810089](#)
- [15] Roberts,M.D.
The Rotation and Shear of a String.
*Class.Q.Grav.***20**(2003)507-519. [hep-th/0204236](#)
- [16] Roberts,M.D.
The Relative Motion of Membranes.
[gr-qc/0404094](#)
- [17] Tseytlin,A.A.
Sigma Model Approach to String Theory.
*Math.Rev.**91c:81117* *Int.J.Mod.Phys.***A4**(1989)1257.